


Longitudinal study of antimicrobial residue in broiler meat in Vietnamese and Bangladeshi farms

Pelligand L¹, Seow K², Butt S¹, Thuy HT⁴, Huong LQ³, Son DT³, Ngoc PT³, Hoa PTT⁴, Conan A⁶, Hasan M⁸, Samad M.A⁸, Roy R.K⁹, Uddin M.H⁹, Dhar P⁹, Mahmud R⁹, Islam S.K.M.A⁹, Biswas P.K⁹, Hoque M.A⁹, Cheah YC², Zhang P², Wang Y², Blake D¹, Hurtaud-Pessel D⁵, Conway P², Fournié G¹, Tomley F¹



Most frequent residue Tilmicosin:
12 Bangladesh 2 Vietnam
Long Withdrawal period 12 days

1 Royal Veterinary College, London, UK, 2 Nanyang Technical University, Singapore, 3 National Institute of Veterinary Research, Ha Noi, Vietnam, 4 CIRAD(French Agricultural Research Centre for International Development), Ha Noi, Vietnam, 5 ANSES (French Agency for Food, Environmental and Occupational Health & Safety), Fougères, France, 6 Centre for Applied One Health Research and Policy Advice, City University of Hong Kong, Hong Kong SAR, 8 Bangladesh Livestock Research Institute, Savar, Dhaka;, 9 Chattogram Veterinary and Animal Sciences University, Chattogram